



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

196. Proposed by L. E. NEWCOMB, Los Gatos, California.

Find the r th term of $\left(x - \frac{1}{x}\right)^n \equiv z^n$ in terms of z .

I. Solution by G. W. GREENWOOD, B. A. (Oxon), Professor of Mathematics and Astronomy, McKendree College, Lebanon Ill.

Let $x = \cos\theta + i\sin\theta$, then $z = 2i\sin\theta$, and $x^r = \cos r\theta + i\sin r\theta$. This enables us to express any term of $(x - 1/x)^n$ in terms of θ , i. e., in terms of z .

II. Solution by F. D. POSEY, A. B., San Mateo, Cal.; G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va., and GRACE M. BAREIS, Bala, Pa.

$$x - 1/x = z, \therefore x = \frac{z \pm \sqrt{z^2 + 4}}{2}.$$

$$\begin{aligned} \text{The } r\text{th term of } (x - 1/x)^n \text{ is } & \frac{(-1)^{r-1} n(n-1) \dots (n-r+2)}{(r-1)!} x^{n-r+1} \left(\frac{1}{x}\right)^{r-1} \\ & = \frac{(-1)^{r-1} n(n-1) \dots (n-r+2)}{(r-1)!} x^{n-2r+2} \\ & = \frac{(-1)^{r-1} n(n-1) \dots (n-r+2)}{(r-1)!} \left(\frac{z \pm \sqrt{z^2 + 4}}{2}\right)^{n-2r+2}. * \end{aligned}$$

197. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, O.

$$\text{Solve } (18)^{4(2-x)} = (54\sqrt{2})^{3x-2}.$$

Solution by W. W. LANDIS, Dickinson College, Carlisle, Pa.

Writing the equation in the form $(1) \dots 18^{4(2-x)} = 18^{3(3x-2)}$ we find $x = \frac{2}{17}$. †

Also solved by G. W. Greenwood, J. E. Sanders, A. H. Holmes, F. D. Posey, R. A. Wells, G. I. Hopkins, H. R. Higley, G. B. M. Zerr, E. L. Sherwood, Grace M. Bareis, J. Scheffer, L. E. Newcomb.

GEOMETRY.

219A. Proposed by H. F. MacNEISH, A. B., Assistant in Mathematics, University High School, Chicago, Ill.

Draw through a given point a line which shall divide a given quadrilateral into two equivalent parts: (1) when the point lies in a side of the quadrilateral; (2) when the point is without; (3) within the quadrilateral.

*Dr. Zerr also gives the values

$$\begin{aligned} & (-1)^{r-1} {}_n C_{r-1} (z + 1/z - 1/z^3 + 1/z^5 - \dots)^{n-2r+2}, \\ & (-1)^r {}_n C_{r-1} (1/z - 1/z^3 + 1/z^5 - \dots)^{n-2r+2}. \end{aligned}$$

†Owing to the periodicity of the exponential function a^y may be written $e^{y \log a + 2n\pi i}$ (n = integer). It now follows from (1) that $e^{\frac{1}{2}(22-17x) \log 18 + 2n\pi i} = 1$, whence $\frac{1}{2}(22-17x) \log 18 + 2n\pi i = 0$, and therefore $x = \frac{2}{17} + (4n\pi i / 17 \log 18)$. When $n=0$, $x = \frac{2}{17}$. ED.